

BTK (13U9) Rabbit Monoclonal Antibody

Catalog# UZ-MA00481

Product information

产品货号	UZ-MA00481
别名	AGMX1; Tyrosine-protein kinase BTK; BTK; Agammaglobulinemia tyrosine kinase; ATK; B-cell progenitor kinase; BPK; Bruton tyrosine kinase; ATK; BPK;
产品名称	BTK (13U9) Rabbit Monoclonal Antibody
纯度	Affinity-chromatography
类别	抗体产品
基因名称	BTK
推荐应用	WB,IHC-P,IF-ICC,FCM,IP,IF-P
反应种属	Human,Mouse,Rat
存储缓冲液	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Human Gene ID	695
Human Swissprot No.	Q06187
免疫原	Recombinant protein of human BTK
稀释度	WB 1:1000, IHC-P/IF-P 1:500-1:2000, IF-ICC 1:100, FCM 1:200-1:500, IP 1:20-1:50
参考分子量	76kDa
预测分子量	76kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	IgG
注意事项	BTK Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
细胞定位	Cytoplasm. Cell membrane; Peripheral membrane protein. Nucleus. Note=In steady state, BTK is predominantly cytosolic Following B-cell receptor (BCR) engagement by antigen, translocates to the plasma membrane through its PH domain. Plasma membrane localization is a critical step in the activation of BTK. A fraction of BTK also shuttles between the nucleus and the cytoplasm, and nuclear export is mediated by the nuclear export receptor CRM1
Clonality	Monoclonal