

CCL2 (6C7) Rabbit Monoclonal Antibody

Catalog# UZ-MA00602

Product information

产品货号	UZ-MA00602
别名	HC11; MCAF; MCP1; MCP-1; SCYA2; GDCF-2; SMC-CF; HSMCR30
产品名称	CCL2 (6C7) Rabbit Monoclonal Antibody
类别	抗体产品
推荐应用	WB,IHC-P,IF-ICC,FCM,IP
反应种属	Human
存储缓冲液	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
免疫原	A synthetic peptide of human CCL2
稀释度	WB 1:1000, IHC-P/IF-P 1:1000-1:10000, IF-ICC 1:50, FCM 1:200-1:500, IP 1:20-1:50
参考分子量	13-15kDa
预测分子量	11kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	IgG
背景介绍	This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Chemokines are a superfamily of secreted proteins involved in immunoregulation and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. [provided by RefSeq, Jul 2013]
纯化	Affinity Purification
Uniprot ID	P13500
Clonality	Monoclonal