

PEG10 (7C18) Rabbit Monoclonal Antibody

Catalog# UZ-MA03459

Product information

产品货号	UZ-MA03459
别名	EDR; HB-1; Mar2; RTL2; MEF3L; Mart2; RGAG3; SIRH1
产品名称	PEG10 (7C18) Rabbit Monoclonal Antibody
类别	抗体产品
推荐应用	WB,IF-ICC,FCM
反应种属	Human,Mouse,Rat
存储缓冲液	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
免疫原	A synthetic peptide of human PEG10
稀释度	WB 1:1000-1:5000, IF-ICC 1:100-1:200, FCM 1:50-1:100
参考分子量	100,55kDa
预测分子量	80kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	IgG
背景介绍	This is a paternally expressed imprinted gene that is thought to have been derived from the Ty3/Gypsy family of retrotransposons. It contains two overlapping open reading frames, RF1 and RF2, and expresses two proteins: a shorter, gag-like protein (with a CCHC-type zinc finger domain) from RF1; and a longer, gag/pol-like fusion protein (with an additional aspartic protease motif) from RF1/RF2 by -1 translational frameshifting (-1 FS). While -1 FS has been observed in RNA viruses and transposons in both prokaryotes and eukaryotes, this gene represents the first example of -1 FS in a eukaryotic cellular gene. This gene is highly conserved across mammalian species and retains the heptanucleotide (GGGAAAC) and pseudoknot elements required for -1 FS. It is expressed in adult and embryonic tissues (most notably in placenta) and reported to have a role in cell proliferation, differentiation and apoptosis. Overexpression of this gene has been associated with several malignancies, such as hepatocellular carcinoma and B-cell lymphocytic leukemia. Knockout mice lacking this gene showed early embryonic lethality with placental defects, indicating the importance of this gene in embryonic development. Additional isoforms resulting from alternatively spliced transcript variants, and use of upstream non-AUG (CUG) start codon have been reported for this gene. [provided by RefSeq, Oct 2014]
纯化	Affinity Purification
Uniprot ID	Q86TG7
Clonality	Monoclonal