

POMC (19F3) Rabbit Monoclonal Antibody

Catalog# UZ-MA03600

Product information

产品货号	UZ-MA03600
别名	LPH; MSH; NPP; POC; ACTH; CLIP
产品名称	POMC (19F3) Rabbit Monoclonal Antibody
类别	抗体产品
推荐应用	WB,IHC-P,IP,IF-P
反应种属	Human,Mouse,Rat
存储缓冲液	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
免疫原	A synthetic peptide of human POMC
稀释度	WB 1:1000-1:5000, IHC-P/IF-P 1:1000-1:10000, IP 1:50-1:100
参考分子量	29kDa
预测分子量	29kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	IgG
背景介绍	This gene encodes a preproprotein that undergoes extensive, tissue-specific, post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the preproprotein and, depending on tissue type and the available convertases, processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation. These include several distinct melanotropins, lipotropins, and endorphins that are contained within the adrenocorticotrophin and beta-lipotropin peptides. The antimicrobial melanotropin alpha peptide exhibits antibacterial and antifungal activity. Mutations in this gene have been associated with early onset obesity, adrenal insufficiency, and red hair pigmentation. Alternatively spliced transcript variants encoding the same protein have been described. [provided by RefSeq, Jan 2016]
纯化	Affinity Purification
Uniprot ID	P01189
Clonality	Monoclonal