

SA2 (6W17) Rabbit Monoclonal Antibody

Catalog# UZ-MA04059

Product information

产品货号	UZ-MA04059
别名	SA 2; SA-2; SA2; SCC3 homolog 2; SCC3B; stag2; Stromal antigen 2;
产品名称	SA2 (6W17) Rabbit Monoclonal Antibody
纯度	Affinity-chromatography
类别	抗体产品
基因名称	STAG2
推荐应用	WB,IHC-P,IF-ICC,FCM,IF-P
反应种属	Human,Mouse,Rat
存储缓冲液	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Human Gene ID	10735
Human Swissprot No.	Q8N3U4
免疫原	Recombinant protein of human SA2
稀释度	WB 1:1000, IHC-P/IF-P 1:200-1:2000, IF-ICC 1:200-1:500, FCM 1:100-1:200
参考分子量	141kDa
预测分子量	141kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	IgG
注意事项	SA2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
细胞定位	Nucleus. Chromosome. Chromosome, centromere. Note=Associates with chromatin. Before prophase it is scattered along chromosome arms. During prophase, most of cohesin complexes dissociate from chromatin probably because of phosphorylation by PLK1, except at centromeres, where cohesin complexes remain. At anaphase, the RAD21 subunit of cohesin is cleaved, leading to the dissociation of the complex from chromosomes, allowing chromosome separation. In germ cells, cohesin complex dissociates from chromatin at prophase I, and may be replaced by a meiosis-specific cohesin complex
Clonality	Monoclonal