

Smad2 (Phospho Thr8) /Smad3 (Thr8) (9U7) Rabbit Monoclonal Antibody

Catalog# UZ-MA04383

Product information

产品货号	UZ-MA04383
别名	JV18; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2
产品名称	Smad2 (Phospho Thr8) /Smad3 (Thr8) (9U7) Rabbit Monoclonal Antibody
类别	抗体产品
推荐应用	WB,IF-ICC,IP
反应种属	Human,Mouse,Rat
存储缓冲液	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
免疫原	A synthetic phosphopeptide corresponding to residues surrounding Thr8 of human Smad2
稀释度	WB 1:1000, IF-ICC 1:50, IP 1:20-1:50
参考分子量	58,62kDa
预测分子量	52,48kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	IgG
背景介绍	The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, May 2012]
纯化	Affinity Purification
Uniprot ID	Q15796,P84022
Clonality	Monoclonal