

Acetyl p53 (K386) Rabbit Polyclonal Antibody

Catalog# UZ-PA01202

Product information

产品货号	UZ-PA01202
别名	TP53; P53; Cellular tumor antigen p53; Antigen NY-CO-13; Phosphoprotein p53; Tumor suppressor p53
产品名称	Acetyl p53 (K386) Rabbit Polyclonal Antibody
类别	抗体产品
基因名称	TP53
推荐应用	IHC-P,IF-P,IF-F,IF-ICC,WB,ELISA
反应种属	Human,Rat,Mouse
存储缓冲液	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N
Human Gene ID	7157
Human Swissprot No.	P04637
免疫原	The antiserum was produced against synthesized peptide derived from human p53 around the acetylated site of Lys386. AA range:344-393
稀释度	WB 1:500-2000, IHC-P 1:100-1:300, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:40000. Not yet tested in other applications.
参考分子量	44kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	Rabbit,IgG
背景介绍	tumor protein p53(TP53) Homo sapiens This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons (PMIDs: 12032546, 20937277). [provided by RefSeq, Feb 2013],
细胞定位	Cytoplasm . Nucleus . Nucleus, PML body . Endoplasmic reticulum . Mitochondrion matrix . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Recruited into PML bodies together with CHEK2 (PubMed:12810724). Translocates to mitochondria upon oxidative stress (PubMed:22726440). Translocates to mitochondria in response to mitomycin C treatment (PubMed:27323408). .; [Isoform 1]: Nucleus . Cytoplasm. Predominantly nuclear but localizes to the cytoplasm when expressed with isoform 4.; [Isoform 2]: Nucleus. Cytoplasm. Localized mainly in the nucleus with minor staining in the cytoplasm.; [Isoform 3]: Nucleus. Cytoplasm. Localized in the nucleus in most cells but found in the cytoplasm in some cells.; [Isoform 4]: Nucleus. Cytoplasm. Predominantly nuclear but translocates to the cytoplasm following cell stress.; [Isoform 7]: Nucleus. Cytoplasm. Localized mainly in the nucleus with minor staining in the cytoplasm.; [Isoform 8]: Nucleus. Cytoplasm. Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli.; [Isoform 9]: Cytoplasm.
纯化	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Acetyl p53 (K386) Rabbit Polyclonal Antibody

Catalog# UZ-PA01202 (cont.)

Product information

Clonality

Polyclonal