

Btk (Phospho Ser180) Rabbit Polyclonal Antibody

Catalog# UZ-PA01766

Product information

产品货号	UZ-PA01766
别名	Tyrosine-protein kinase BTK (EC 2.7.10.2) (Agammaglobulinaemia tyrosine kinase) (ATK) (B-cell progenitor kinase) (BPK) (Bruton tyrosine kinase);AGMX1;ATK;BPK
产品名称	Btk (Phospho Ser180) Rabbit Polyclonal Antibody
类别	抗体产品
基因名称	BTK
推荐应用	WB
反应种属	Human,Rat,Mouse
存储缓冲液	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N .
Human Gene ID	695
Human Swissprot No.	Q06187
免疫原	Synthesized phosho peptide around human Btk (Ser180)
稀释度	WB 1:1000-2000
预测分子量	80kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	Rabbit,IgG
背景介绍	The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2013],
细胞定位	Cytoplasm. Cell membrane; Peripheral membrane protein. Nucleus. In steady state, BTK is predominantly cytosolic. Following B-cell receptor (BCR) engagement by antigen, translocates to the plasma membrane through its PH domain. Plasma membrane localization is a critical step in the activation of BTK. A fraction of BTK also shuttles between the nucleus and the cytoplasm, and nuclear export is mediated by the nuclear export receptor CRM1.
纯化	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Clonality	Polyclonal