

CBLB Rabbit Polyclonal Antibody

Catalog# UZ-PA02141

Product information

产品货号	UZ-PA02141
别名	RNF56;Nbla00127
产品名称	CBLB Rabbit Polyclonal Antibody
类别	抗体产品
基因名称	CBLB
推荐应用	WB,ELISA
反应种属	Human,Rat,Mouse
存储缓冲液	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Human Gene ID	868
Human Swissprot No.	Q13191
免疫原	Synthesized peptide derived from part region of human protein
稀释度	WB 1:500-2000, ELISA 1:5000-20000
预测分子量	108kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	Rabbit,IgG

CBLB Rabbit Polyclonal Antibody

Catalog# UZ-PA02141 (cont.)

Product information

背景介绍

domain:The N-terminus is composed of the phosphotyrosine binding (PTB) domain, a short linker region and the RING-type zinc finger. The PTB domain, which is also called TKB (tyrosine kinase binding) domain, is composed of three different subdomains: a four-helix bundle (4H), a calcium-binding EF hand and a divergent SH2 domain. domain:The RING-type zinc finger domain mediates binding to an E2 ubiquitin-conjugating enzyme. domain:The UBA domain interacts with poly-ubiquitinated proteins. function:E3 ubiquitin-protein ligase which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and transfers it to substrates, generally promoting their degradation by the proteasome. Negatively regulates TCR (T-cell receptor), BCR (B-cell receptor) and FCER1 (high affinity immunoglobulin epsilon receptor) signal transduction pathways. In naive T-cells, inhibits VAV1 activation upon TCR engagement and imposes a requirement for CD28 costimulation for proliferation and IL-2 production. Also acts by promoting PIK3R1/p85 ubiquitination, which impairs its recruitment to the TCR and subsequent activation. In activated T-cells, inhibits PLCG1 activation and calcium mobilization upon restimulation and promotes anergy. In B-cells, acts by ubiquitinating SYK and promoting its proteasomal degradation. May also be involved in EGFR ubiquitination and internalization. miscellaneous:This protein has one functional calcium-binding site. pathway:Protein modification; protein ubiquitination. PTM:Auto-ubiquitinated upon EGF-mediated cell activation or upon T-cell costimulation by CD28; which promotes proteasomal degradation. PTM:Phosphorylated on tyrosine residues upon TCR or BCR activation, and upon various types of cell stimulation. sequence caution:Translated as Arg. similarity:Contains 1 CBL N-terminal domain. similarity:Contains 1 RING-type zinc finger. similarity:Contains 1 SH2 domain. similarity:Contains 1 UBA domain. similarity:Contains 2 EF-hand-like domains. subcellular location:Upon EGF stimulation, associates with endocytic vesicles. subunit:Interacts with SH3 domain-containing proteins LCK, CRK and SORBS1. Interacts with LCP2 and ZAP70. May interact with CBL (By similarity). Interacts with SH3 domain-containing proteins VAV1, FYN, FGR, PLCG1, GRB2, CRKL, PIK3R1 and SH3KBP1/CIN85. Identified in heterotrimeric complexes with SH3KBP1/CIN85, CD2AP and ARHGEF7, where one CBLB peptide binds two copies of the other protein. Interacts with poly-ubiquitinated proteins. Dimerization is required for the binding of poly-ubiquitin, but not for the binding of mono-ubiquitin. tissue specificity:Expressed in placenta, heart, lung, kidney, spleen, ovary and testis, as well as fetal brain and liver and hematopoietic cell lines, but not in adult brain, liver, pancreas, salivary gland, or skeletal muscle. Present in lymphocytes (at protein level).

细胞定位 纯化

Cytoplasm . Upon EGF stimulation, associates with endocytic vesicles.

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Clonality

Polyclonal