

CBP 35 Rabbit Polyclonal Antibody

Catalog# UZ-PA02145

Product information

产品货号	UZ-PA02145
别名	LGALS3; MAC2; Galectin-3; Gal-3; 35 kDa lectin; Carbohydrate-binding protein 35; CBP 35; Galactose-specific lectin 3; Galactoside-binding protein; GALBP; IgE-binding protein; L-31; Laminin-binding protein; Lectin L-29; Mac-2 antigen
产品名称	CBP 35 Rabbit Polyclonal Antibody
类别	抗体产品
基因名称	LGALS3
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N .
Human Gene ID	3958
Human Swissprot No.	P17931
免疫原	The antiserum was produced against synthesized peptide derived from human Galectin 3. AA range:141-190
稀释度	WB 1:500-1:2000, IHC-P 1:100-1:300, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:20000 .Not yet tested in other applications.
预测分子量	33kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	Rabbit,IgG
背景介绍	This gene encodes a member of the galectin family of carbohydrate binding proteins. Members of this protein family have an affinity for beta-galactosides. The encoded protein is characterized by an N-terminal proline-rich tandem repeat domain and a single C-terminal carbohydrate recognition domain. This protein can self-associate through the N-terminal domain allowing it to bind to multivalent saccharide ligands. This protein localizes to the extracellular matrix, the cytoplasm and the nucleus. This protein plays a role in numerous cellular functions including apoptosis, innate immunity, cell adhesion and T-cell regulation. The protein exhibits antimicrobial activity against bacteria and fungi. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Oct 2014],
细胞定位	Cytoplasm . Nucleus. Secreted . Secreted by a non-classical secretory pathway and associates with the cell surface. Can be secreted; the secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10; it results in protein translocation from the cytoplasm into the ERGIC (endoplasmic reticulum-Golgi intermediate compartment) followed by vesicle entry and secretion (PubMed:32272059). .
纯化	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal