

CCL18 Rabbit Polyclonal Antibody

Catalog# UZ-PA02287

Product information

产品货号	UZ-PA02287
别名	C-C motif chemokine 18 (Alternative macrophage activation-associated CC chemokine 1;AMAC-1;CC chemokine PARC;Dendritic cell chemokine 1;DC-CK1;Macrophage inflammatory protein 4;MIP-4;Pulmonary and activation-regulated chemokine;Small-inducible cytokine A18) [Cleaved into: CCL18(1-68); CCL18(3-69); CCL18(4-69)];AMAC1;DCC K1;MIP4;PARC;SCYA18;
产品名称	CCL18 Rabbit Polyclonal Antibody
类别	抗体产品
基因名称	CCL18
推荐应用	IHC-P,ELISA
反应种属	Human,Rat,Mouse
存储缓冲液	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N .
Human Gene ID	6362
Human Swissprot No.	P55774
免疫原	Synthesized peptide derived from human CCL18 AA range: 1-80
稀释度	IHC-P 1:50-200, ELISA (peptide)1:5000-20000
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	Rabbit,IgG
背景介绍	This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for naive T cells, CD4+ and CD8+ T cells and nonactivated lymphocytes, but not for monocytes or granulocytes. This chemokine attracts naive T lymphocytes toward dendritic cells and activated macrophages in lymph nodes. It may play a role in both humoral and cell-mediated immunity responses. [provided by RefSeq, Sep 2014],
细胞定位	Secreted.
纯化	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal