

PDPK1 (Phospho Ser241) (3O2) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00586

Product information

产品货号	HZ-YMA00586
别名	PDPK1;PDK1;3-phosphoinositide-dependent protein kinase 1;hPDK1
产品名称	PDPK1 (Phospho Ser241) (3O2) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	PDPK1
推荐应用	WB,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	5170
Human Swissprot No.	O15530
稀释度	WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000, IP 1:50-1:200
参考分子量	63kDa
预测分子量	63kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Phosphorylates and activates not only PKB/AKT, but also PKA, PKC-zeta, RPS6KA1 and RPS6KB1. May play a general role in signaling processes and in development (By similarity). Isoform 3 is catalytically inactive.,PTM:Phosphorylated on tyrosine and serine/threonine. Phosphorylation on Ser-241 in the activation loop is required for full activity. PDK1 itself can autophosphorylate Ser-241, leading to its own activation.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PDK1 subfamily.,similarity:Contains 1 PH domain.,similarity:Contains 1 protein kinase domain.,subcellular location:Membrane-associated after cell stimulation leading to its translocation. Tyrosine phosphorylation seems to occur only at the plasma membrane.,subunit:Interacts with TUSC4.,tissue specificity:Appears to be expressed ubiquitously.,
细胞定位	Cytoplasm. Nucleus. Cell membrane; Peripheral membrane protein. Cell junction, focal adhesion. Tyrosine phosphorylation seems to occur only at the cell membrane. Translocates to the cell membrane following insulin stimulation by a mechanism that involves binding to GRB14 and INSR. SRC and HSP90 promote its localization to the cell membrane. Its nuclear localization is dependent on its association with PTPN6 and its phosphorylation at Ser-396. Restricted to the nucleus in neuronal cells while in non-neuronal cells it is found in the cytoplasm. The Ser-241 phosphorylated form is distributed along the perinuclear region in neuronal cells while in non-neuronal cells it is found in both the nucleus and the cytoplasm. IGF1 transiently increases phosphorylation at Ser-241 of neuronal PDPK1, resulting in its translocation to other cellular compartments. The tyrosine-phosphorylated form colocalizes with PTK2B in focal adhesions after angiotensin II stimulation.
纯化	Protein A
Clonality	Monoclonal