

Tau (Phospho Ser396) (4U10) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00649

Product information

产品货号	HZ-YMA00649
别名	MAPT;MAPTL;MTBT1;TAU;Microtubule-associated protein tau;Neurofibrillary tangle protein;Paired helical filament-tau;PHF-tau
产品名称	Tau (Phospho Ser396) (4U10) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	MAPT
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	4137
Human Swissprot No.	P10636
稀释度	IHC-P 1:200-1:1000, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000
参考分子量	55-130kDa
预测分子量	79kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	This gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy. [provided by RefSeq, Jul 2008],
细胞定位	Cytoplasm, cytosol . Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Secreted . Mostly found in the axons of neurons, in the cytosol and in association with plasma membrane components (PubMed:10747907). Can be secreted; the secretion is dependent on protein unfolding and facilitated by the cargo receptor TMD10; it results in protein translocation from the cytoplasm into the ERGIC (endoplasmic reticulum-Golgi intermediate compartment) followed by vesicle entry and secretion (PubMed:32272059). .
纯化	Protein A
Clonality	Monoclonal