

cIAP2 (6X5) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00698

Product information

产品货号	HZ-YMA00698
别名	AIP 1;AIP1;API 2;API2;API2;Apoptosis inhibitor 2;Baculoviral IAP repeat containing protein 3;Baculoviral IAP repeat containing protein 3;Baculoviral IAP repeat-containing protein 3;BIRC 3;BIRC3;BIRC3;BIRC3_HUMAN;C IAP2;C-IAP2;CIAP 2;CIAP 2;CIAP2;HAIP 1;HAIP1;HAIP1;HIAP 1;HIAP-1;HIAP1;IAP homolog C;IAP-1;Inhibitor of apoptosis protein 1;Inhibitor of apoptosis protein 1;MALT 2;MALT2;Mammalian IAP homolog C;MIHC;MIHC;RING finger protein 49;RNF49;TNFR2 TRAF signaling complex protein 1;TNFR2 TRAF signalling complex protein;TNFR2-TRAF-signaling complex protein 1.
产品名称	cIAP2 (6X5) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	BIRC3
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	330
Human Swissprot No.	Q13489
稀释度	IHC-P 1:200-1:500, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:500-1:20000, IP 1:50-1:200
参考分子量	68kDa
预测分子量	68kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus IAP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq, Aug 2011],
细胞定位	Cytoplasm . Nucleus .
纯化	Protein A
Clonality	Monoclonal