

PPAR- γ (19R7) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00774

Product information

产品货号	HZ-YMA00774
别名	PPARG;NR1C3;Peroxisome proliferator-activated receptor gamma;PPAR-gamma;Nuclear receptor subfamily 1 group C member 3
产品名称	PPAR- γ (19R7) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	PPARG
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	5468
Human Swissprot No.	P37231
稀释度	IHC-P 1:500-1:1000, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:500-1:20000, IP 1:50-1:200,
参考分子量	53kDa
预测分子量	58kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	peroxisome proliferator activated receptor gamma(PPARG) Homo sapiens This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) subfamily of nuclear receptors. PPARs form heterodimers with retinoid X receptors (RXRs) and these heterodimers regulate transcription of various genes. Three subtypes of PPARs are known: PPAR-alpha, PPAR-delta, and PPAR-gamma. The protein encoded by this gene is PPAR-gamma and is a regulator of adipocyte differentiation. Additionally, PPAR-gamma has been implicated in the pathology of numerous diseases including obesity, diabetes, atherosclerosis and cancer. Alternatively spliced transcript variants that encode different isoforms have been described. [provided by RefSeq, Jul 2008],
细胞定位	Nucleus. Cytoplasm. Redistributed from the nucleus to the cytosol through a MAP2K1/MEK1-dependent manner. NOCT enhances its nuclear translocation.
纯化	Protein A
Clonality	Monoclonal