

Rpb1 (Phospho Ser2) (17Z17) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00786

Product information

产品货号	HZ-YMA00786
别名	DNA-directed RNA polymerase II subunit RPB1;RNA polymerase II subunit B1;DNA-directed RNA polymerase II subunit A;DNA-directed RNA polymerase III largest subunit;RNA-directed RNA polymerase II subunit RPB1;POLR2
产品名称	Rpb1 (Phospho Ser2) (17Z17) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	POLR2A
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	5430
Human Swissprot No.	P24928
稀释度	IHC-P 1:100-1:500, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:500-1:20000, IP 1:50-1:200,
参考分子量	270kDa
预测分子量	217kDa
运输及保存条件	Biological ice bag transportation. Store at -20℃ for at least 12 months(Do not lower than -25℃). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	This gene encodes the largest subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. The product of this gene contains a carboxy terminal domain composed of heptapeptide repeats that are essential for polymerase activity. These repeats contain serine and threonine residues that are phosphorylated in actively transcribing RNA polymerase. In addition, this subunit, in combination with several other polymerase subunits, forms the DNA binding domain of the polymerase, a groove in which the DNA template is transcribed into RNA. [provided by RefSeq, Jul 2008],
细胞定位	Nucleus . Cytoplasm . Chromosome . Hypophosphorylated form is mainly found in the cytoplasm, while the hyperphosphorylated and active form is nuclear (PubMed:26566685). Co-localizes with kinase SRPK2 and helicase DDX23 at chromatin loci where unscheduled R-loops form (PubMed:28076779). .
纯化	Protein A
Clonality	Monoclonal