

Thioredoxin (10I12) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00800

Product information

产品货号	HZ-YMA00800
别名	Thioredoxin;Trx;ATL-derived factor;ADF;Surface-associated sulphhydryl protein;SASP;TRDX TRX TRX1
产品名称	Thioredoxin (10I12) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	TXN
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	7295
Human Swissprot No.	P10599
稀释度	IHC-P 1:500-1:2000, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000
参考分子量	12kDa
预测分子量	12kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	The protein encoded by this gene acts as a homodimer and is involved in many redox reactions. The encoded protein is active in the reversible S-nitrosylation of cysteines in certain proteins, which is part of the response to intracellular nitric oxide. This protein is found in the cytoplasm. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011],
细胞定位	Nucleus . Cytoplasm . Secreted . Translocates from the cytoplasm into the nucleus after phorbol 12-myristate 13-acetate induction (PMA) (PubMed:9108029). Predominantly in the cytoplasm in non irradiated cells (PubMed:11118054). Radiation induces translocation of TRX from the cytoplasm to the nucleus (PubMed:11118054). Secreted by a leaderless secretory pathway (PubMed:1332947). .
纯化	Protein A
Clonality	Monoclonal