

PI3-Kinase p110 γ (19X4) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00815

Product information

产品货号	HZ-YMA00815
别名	PIK3CG;Phosphatidylinositol 4;5-bisphosphate 3-kinase catalytic subunit gamma isoform;PI3-kinase subunit gamma;PI3K-gamma;PI3Kgamma;PtdIns-3-kinase subunit gamma;Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit
产品名称	PI3-Kinase p110 γ (19X4) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	PIK3CG
推荐应用	WB,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	5294
Human Swissprot No.	P48736
稀释度	WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000, IP 1:50-1:200,
参考分子量	127kDa
预测分子量	127kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I catalytic subunit of PI3K. Like other class I catalytic subunits (p110-alpha p110-beta, and p110-delta), the encoded protein binds a p85 regulatory subunit to form PI3K. This gene is located in a commonly deleted segment of chromosome 7 previously identified in myeloid leukemias. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jun 2015].
细胞定位	Cytoplasm . Cell membrane .
纯化	Protein A
Clonality	Monoclonal