

PARK7 (18Q11) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00847

Product information

产品货号	HZ-YMA00847
别名	PARK7;Protein DJ-1;Oncogene DJ1;Parkinson disease protein 7
产品名称	PARK7 (18Q11) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	PARK7
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	11315
Human Swissprot No.	Q99497
稀释度	IHC-P 1:4000-1:20000, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000, IP 1:50-1:200,
参考分子量	22kDa
预测分子量	20kDa
运输及保存条件	Biological ice bag transportation. Store at -20℃ for at least 12 months(Do not lower than -25℃). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	The product of this gene belongs to the peptidase C56 family of proteins. It acts as a positive regulator of androgen receptor-dependent transcription. It may also function as a redox-sensitive chaperone, as a sensor for oxidative stress, and it apparently protects neurons against oxidative stress and cell death. Defects in this gene are the cause of autosomal recessive early-onset Parkinson disease 7. Two transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq, Jul 2008],
细胞定位	Cell membrane ; Lipid-anchor . Cytoplasm . Nucleus . Membrane raft . Mitochondrion . Endoplasmic reticulum . Under normal conditions, located predominantly in the cytoplasm and, to a lesser extent, in the nucleus and mitochondrion. Translocates to the mitochondrion and subsequently to the nucleus in response to oxidative stress and exerts an increased cytoprotective effect against oxidative damage (PubMed:18711745). Detected in tau inclusions in brains from neurodegenerative disease patients (PubMed:14705119). Membrane raft localization in astrocytes and neuronal cells requires palmitoylation. .
纯化	Protein A
Clonality	Monoclonal