

NF-κB p65 (Acetyl Lys310) (14H14) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00915

Product information

产品货号	HZ-YMA00915
别名	RELA;NFKB3;Transcription factor p65;Nuclear factor NF-kappa-B p65 subunit;Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3
产品名称	NF-κB p65 (Acetyl Lys310) (14H14) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	RELA
推荐应用	WB,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	5970
Human Swissprot No.	Q04206
稀释度	WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000, IP 1:50-1:200
参考分子量	65kDa
预测分子量	60kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011],
细胞定位	Nucleus . Cytoplasm . Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor (I-kappa-B) (PubMed:1493333). Colocalized with DDX1 in the nucleus upon TNF-alpha induction (PubMed:19058135). Colocalizes with GFI1 in the nucleus after LPS stimulation (PubMed:20547752). Translocation to the nucleus is impaired in L.monocytogenes infection (PubMed:20855622). .
纯化	Protein A
Clonality	Monoclonal