

HDAC3 (2V18) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00992

Product information

产品货号	HZ-YMA00992
别名	HDAC3;Histone deacetylase 3;HD3;RPD3-2;SMAP45
产品名称	HDAC3 (2V18) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	HDAC3
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	8841
Human Swissprot No.	O15379
稀释度	IHC-P 1:200-1:1000, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000, IP 1:50-1:200
参考分子量	49kDa
预测分子量	49kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family. It has histone deacetylase activity and represses transcription when tethered to a promoter. It may participate in the regulation of transcription through its binding with the zinc-finger transcription factor YY1. This protein can also down-regulate p53 function and thus modulate cell growth and apoptosis. This gene is regarded as a potential tumor suppressor gene. [provided by RefSeq, Jul 2008].
细胞定位	Nucleus . Cytoplasm . Cytoplasm, cytosol . Colocalizes with XBP1 and AKT1 in the cytoplasm (PubMed:25190803). Predominantly expressed in the nucleus in the presence of CCAR2 (PubMed:21030595). .
纯化	Protein A
Clonality	Monoclonal