

LIMK-2 (12T16) Rabbit Monoclonal Antibody

Catalog# HZ-YMA01123

Product information

产品货号	HZ-YMA01123
别名	LIMK2;LIM domain kinase 2;LIMK-2
产品名称	LIMK-2 (12T16) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	LIMK2
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	3985
Human Swissprot No.	P53671
稀释度	IHC-P 1:200-1:1000, WB 1:2000-1:10000, IF 1:200-1:1000, ELISA 1:5000-1:20000, IP 1:50-1:200
参考分子量	72kDa
预测分子量	72kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],
细胞定位	Cytoplasm, cytoskeleton, spindle . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .; [Isoform LIMK2a]: Cytoplasm . Nucleus .; [Isoform LIMK2b]: Cytoplasm . Cytoplasm, perinuclear region . Nucleus . Mainly present in the cytoplasm and is scarcely translocated to the nucleus. .
Clonality	Monoclonal