

WTAP (9X5) Rabbit Monoclonal Antibody

Catalog# HZ-YMA01263

Product information

产品货号	HZ-YMA01263
别名	WTAP; KIAA0105; Pre-mRNA-splicing regulator WTAP; Female-lethal; 2; D homolog; hFL; 2; D; WT1-associated protein; Wilms tumor 1-associating protein
产品名称	WTAP (9X5) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	WTAP
推荐应用	WB, IHC-P, IF-P, IF-F, IF-ICC, IP, ELISA
反应种属	Human
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA
Human Gene ID	9589
Human Swissprot No.	Q15007
稀释度	IHC-P 1:200-1:1000; WB 1:2000-1:10000; IF-P/IF-F/IF-ICC 1:200-1:1000; ELISA 1:5000-1:20000; IP 1:50-1:200;
参考分子量	55kDa
预测分子量	44kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months (Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG, Kappa
背景介绍	The Wilms tumor suppressor gene WT1 appears to play a role in both transcriptional and posttranscriptional regulation of certain cellular genes. This gene encodes a WT1-associating protein, which is a ubiquitously expressed nuclear protein. Like WT1 protein, this protein is localized throughout the nucleoplasm as well as in speckles and partially colocalizes with splicing factors. Alternative splicing of this gene results in several transcript variants encoding three different isoforms. [provided by RefSeq, Jul 2012],
细胞定位	Nucleus speckle . Nucleus, nucleoplasm . Cytoplasm . Mainly nuclear with some fraction located in the cytoplasm. ZC3H13 is required to anchor component of the MACOM subcomplex, such as VIRMA, in the nucleus. .
Clonality	Monoclonal