

CD79b (14I19) Rabbit Monoclonal Antibody

Catalog# HZ-YMA01265

Product information

产品货号	HZ-YMA01265
别名	CD79B;B29;IGB;B-cell antigen receptor complex-associated protein beta chain;B-cell-specific glycoprotein B29;Ig-beta;Immunoglobulin-associated B29 protein;CD79b
产品名称	CD79b (14I19) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	CD79B
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	974
Human Swissprot No.	P40259
稀释度	IHC-P 1:200-1:1000;WB 1:2000-1:10000;IF-P/IF-F/IF-ICC 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-1:200;
参考分子量	32-36kDa
预测分子量	26kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the Ig-beta protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008],
细胞定位	Cell membrane; Single-pass type I membrane protein. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts. .
Clonality	Monoclonal