

NCoA 3 (17D17) Rabbit Monoclonal Antibody

Catalog# HZ-YMA01363

Product information

产品货号	HZ-YMA01363
别名	NCOA3;AIB1;BHLHE42;RAC3;TRAM1;Nuclear receptor coactivator 3;NCoA-3;ACTR;Amplified in breast cancer 1 protein;AIB-1;CBP-interacting protein;pCIP;Class E basic helix-loop-helix protein 42;bHLHe42;Receptor-associated coactivator 3;RAC-3;Steroid receptor coactivator protein 3;SRC-3;Thyroid hormone receptor activator molecule 1;TRAM-1
产品名称	NCoA 3 (17D17) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	NCOA3
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	8202
Human Swissprot No.	Q9Y6Q9
稀释度	IHC-P 1:200-1:500;WB 1:2000-1:10000;IF-P/IF-F/IF-ICC 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-1:200;
参考分子量	160kDa
预测分子量	155kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Several transcript variants encoding different isoforms have been found for this gene. In addition, a polymorphic repeat region is found in the C-terminus of the encoded protein. [provided by RefSeq, Mar 2010],
细胞定位	Cytoplasm. Nucleus. Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus.
Clonality	Monoclonal