

PI 3-Kinase p110δ (16T15) Rabbit Monoclonal Antibody

Catalog# HZ-YMA01540

Product information

产品货号	HZ-YMA01540
别名	PIK3CD;Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform;PI3-kinase subunit delta;PI3K-delta;PI3Kdelta;PtdIns-3-kinase subunit delta;Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit delta;PtdIns-3-kinase subunit p110-delta;p110delta
产品名称	PI 3-Kinase p110δ (16T15) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	PIK3CD
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
稀释度	IHC-P 1:200-1:1000, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000
参考分子量	119kDa
预测分子量	119kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
细胞定位	In humans, the highest levels of expression are seen in peripheral blood mononuclear cells, spleen, and thymus, and low levels of expression in testes, uterus, colon, and small intestine but not in other tissues examined including prostate, heart, brain, and liver (PubMed:9235916). Isoform 2 is expressed in normal thymus, lung and spleen tissues, and is detected at low levels in normal lysates from colon and ovarian biopsies, at elevated levels in lysates from colorectal tumors and is abundantly expressed in some ovarian tumors (at protein level). Both isoform 1 and isoform 2 are widely expressed. Isoform 1 is expressed predominantly in leukocytes.
Clonality	Monoclonal