

Bcl-10 (19R16) Rabbit Monoclonal Antibody

Catalog# HZ-YMA01576

Product information

产品货号	HZ-YMA01576
别名	BCL10;CIPER;CLAP;B-cell lymphoma/leukemia 10;B-cell CLL/lymphoma 10;Bcl-10;CARD-containing molecule enhancing NF-kappa-B;CARD-like apoptotic protein;hCLAP;CED-3/ICH-1 prodomain homologous E10-like regulator;CIPER;Cellular homolog
产品名称	Bcl-10 (19R16) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	BCL10
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	8915
Human Swissprot No.	O95999
稀释度	IHC-P 1:200-1:1000;WB 1:2000-1:10000;IF-P/IF-F/IF-ICC 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-1:200;
参考分子量	32kDa
预测分子量	26kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	IgG,Kappa
背景介绍	This gene was identified by its translocation in a case of mucosa-associated lymphoid tissue (MALT) lymphoma. The protein encoded by this gene contains a caspase recruitment domain (CARD), and has been shown to induce apoptosis and to activate NF-kappaB. This protein is reported to interact with other CARD domain containing proteins including CARD9, 10, 11 and 14, which are thought to function as upstream regulators in NF-kappaB signaling. This protein is found to form a complex with MALT1, a protein encoded by another gene known to be translocated in MALT lymphoma. MALT1 and this protein are thought to synergize in the activation of NF-kappaB, and the deregulation of either of them may contribute to the same pathologic process that leads to the malignancy. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],
细胞定位	Cytoplasm, perinuclear region . Membrane raft . Appears to have a perinuclear, compact and filamentous pattern of expression. Also found in the nucleus of several types of tumor cells. Colocalized with DPP4 in membrane rafts. .
纯化	Protein A
Clonality	Monoclonal